

The barley α -thionin promoter is rich in negative regulatory motifs and directs tissue-specific expression of a reporter gene in tobacco

José A. Fernández, Manuel Moreno, María J. Carmona, Atilio Castagnaro
and Francisco García-Olmedo

Laboratorio de Bioquímica y Biología Molecular, E.T.S. Ingenieros Agrónomos-UPM, Madrid (Spain)

Key words: Negative regulatory element; Promoter analysis; Thionin; Tissue-specificity

The promoter of the barley α -thionin gene (1.6 kb) fused to the β -glucuronidase (GUS) gene directs temporally-controlled, tissue-specific expression in the endosperm of transgenic tobacco. The nucleotide sequence of this promoter shows negative regulatory motifs which have been functionally analyzed in other genes.

Thionins are cysteine-rich 5 kDa plant polypeptides that have been identified in a limited number of wide-ranging taxa and are toxic to plant pathogens (see Refs. 1,2 for reviews). Known thionin sequences have been classified into five structural types, three of which are found in wheat and barley (I,II,V), where types I and V are very abundant in endosperm and type II in leaves [2]. Genes encoding type I [3–5] and type V [6] are expressed during the cell-proliferation phase of endosperm development (first half), while those of type II are expressed in etiolated leaves and induced under certain stress conditions, including infection by pathogens [7–9]. While the number of type II genes has been estimated at 10–100 copies per haploid genome [7,9], only 1–2 copies per haploid genome seem to be present for the other two types [5,6,10]. This implies that high accumulation of thionins, which occurs to a similar extent in both cases, is achieved by the number of gene copies in the first case and by strong promoters in the second.

We have now sequenced the promoter of the α -thionin gene from barley (type I) and have found that this promoter directs tissue-specific, temporally-con-

trolled expression of the β -glucuronidase (GUS) reporter gene in transgenic tobacco.

A DNA fragment (*Xba*I-*Nco*I; –1.6 kb to +45 b) upstream of the coding region of the barley α -thionin [5] was subcloned and both strands completely sequenced, using standard protocols for the dideoxy chain termination method. The sequence is shown in Fig. 1 and the main putative regulatory features are summarized in Table I. The nucleotide sequence contains several motifs that are identical or very similar to sequences that have been functionally characterized as negative regulatory elements in other plant genes.

A gene fusion involving the α -thionin promoter and the bacterial β -glucuronidase (GUS) reporter gene was carried out as indicated in Fig. 2A. This gene construction (α -TH-GUS) was introduced and expressed in tobacco using the Bin19 binary vector system [11]. Non-transformed tobacco and transformants carrying a CaMV-35S-GUS fusion (hereafter 35S) were used as negative and positive controls, respectively.

GUS activity was measured fluorimetrically [12] in seeds, leaves, stems, and roots from individual regenerants and the data are summarized in Fig. 2B. Expression levels well above those of the 35S positive control were observed in seeds, while only background fluorescence was observed in the other parts of the plant analyzed. Developing seeds were collected at different days after pollination and endosperms were separated from embryos and coats by hand dissection. GUS activity in α -TH-GUS seeds was restricted to the en-

-1654 TCTAGAGCATTTAAATTTAGCTAAAGTAGGGTGAGGATGAAAGGATGAAGTAGCTAACCTTTTAGAACACTTGTGTAGTTGCTGCACCACCAAAACCTA
-1555 GACAAATCTTGGGAAATGGAGCTTGGAGGTCGAGCTTGGAGAGGAGAAAGCTTAAGTGTGGCTCGGGCATTTTCATCGAACACCTCATGTGCATGCAT
-1456 AGAACGGTGAACATAGCAGGGCATGCACACCTCCACACGGCCAAAAACAGAGGAGAGGGTGGGCGAGGGGATATATATAGTATCTCTTTGGT
-1357 CCCGGTTGGTGGCCAGAACGGGACAGAACTGACTTTTAGTCCCGGTTCCACCCACCAACCGGGACTAAAGGTGCTGCACGAGGAGTGAGGACCATT
-1258 GGTCTCGGTTCTGTCATGGAACAGGACCAAGGGGTGAGACGAACCGAGACCTGCCCCAGCCGACGCCCTGGCCTCATGAACAGGGAGCTGATATGG
-1159 TCCCGGTTCTGAACAGAATCGGGATTAAAGCTCTTATCTCGCCTCGACCAAGGCCCTGTTTTCTACTAGTGTGATTTTCTGCTCAGACACATGCCACT
-1060 CCTTCTCTTCCCACGTCAATTTCTGAAGATCAATTTCTGCTGACTCATGAGTGAATCTAGCATCGATACGACATCTTATGTTAGACATTTT
-961 TATATGATGATTTTATTAATGAGGATCTTAAGATACCATAGTTTGTGCTTCGTTCCGATTCGAGAGGTTTTTGGATCCACCCAGGAGGATTAGA
-862 CAAGGTGCTAAGTGCATCGAAGGACTTATGAACAACTCACATATGCATGTATATACTTAGCCTTTTCTAGCTAAGATTTCTCCATTTTATGGTTTA
-763 GCAACAAATTCGCATGCATATGATTTTTCATAGATTACTCGTTTGACATTGTTGTTTCTTCCTGGGTCTCCATGCTATCTCTTCTTAATTA
-664 ATTACTTTGCTACGTTAACTTTTCCGATGTGACATTTTTCAGCAGTTGAGCGCTGAAAAGGCCCTAAAGCACATAAGGACAAGAACTAGCTAACTG
-565 CCCAAAAGTTTCGGTTTTGAGGGATGCATTACCAACACATAAGTGATATACCATGTGAACGTGGTTCGCAGATGGTTTGATACAAACATGTTTGCTTGC
-466 ACGTATTTTATAACCATCATCTTTATCATACATAACCAAAATTTCTCTTGGGATGGATGACAGCCATAAGGTCGTCTTTTTCCATTGCCGCTGTGA
-367 TTAAGACACTTTGTTTTCATTGCCAAAGTCGGCTCCTTGGCAAGGCAAGAGGACAAGTACAATTTTGTGATAAGTCTCTTTTAAAGTATCATG
-268 TTTAAGGAGTATATGCAACTAGAAACCGATTTTGTCTATGTCGATGTGGTTAATGATTGTGAGTGGAGTTAGAAGTGGAGTGAAGTAGAGTTTGA
-169 GGTGAGTGTAGCACTTCAACACACTCTGTAAGTAAACAGCTAAGAGGTTTTGTTTTTCGTCGGCTGGCATGAACACGTAACATTTAACCTACAACAAC
-70 ATCGCTTGGACATCCTGTGTGCCACAACCACCATCTCTATACTAAGGCCATCCAAGCCTAGTCTCAG

Fig. 1. Nucleotide sequence of the α -thionin promoter from barley. Boxes with high homology to silencer motifs in other plant promoters are shadowed (see Table I). The following motifs are numbered and over or underlined (stars indicate complementary strands): β -phaseolin positive elements CCACC/A(1) and AACACA(2); yeast GCN4 binding site ATGAC(3); consensus of boxes II and III of rbcS-3A GPuT/AAT/A(4) and box-II core of the same gene GGTTAA (5). A 10 bp direct repeat is indicated by arrows and two hairpin loops by an interrupted overline. A region with homology to *Drosophila* ORF-1 promoter is boxed and the putative TATA box is bold.

TABLE I

Summary of putative regulatory motifs in the α -thionin promoter

| α -Thionin promoter | Other plant promoters | Ref. |
|---|---|------|
| TATATATAAA (-1379, -1371 bp) | Silencer motifs | |
| CATCTTAAGATA (-934, -923 bp) | Cab identical (-964 *, -972 * bp) | 13 |
| TCAAATTTGTG (-1040, -1020 bp) | (-953 *, -968 * bp) | 13 |
| (-1022, -1012 bp) | Cab similar (-1067, -1057 bp) | 13 |
| ATGACTTTGTATTAAAAA (-965, -938 bp) | rbcS-3A similar (-160, -150 bp) | 14 |
| | Cab similar (-1059 *, -1077 * bp) | 13 |
| | rbcS-3A similar (-104 *, -121 * bp) | 14 |
| | CHS 15 similar (-289, -273 bp) | 15 |
| CATTTAAAATTAGCTAACTAGC (-1646, -1623 bp) | CHS 15 similar (-321, -301 bp) | 15 |
| | (-211 *, -237 * bp) | 15 |
| CCACC/A (9 repeats in Fig. 1) | Identity to other motifs | |
| AACACA (2 repeats in Fig. 1) | β -Phs pentamer box (-468, -391 bp) | 16 |
| ATGAC (4 repeats in Fig. 1) | hexamer box (-391, -295 bp) | 16 |
| GPuT/AAAT/A (2 repeats in Fig. 1) | binding consensus yeast GCN4 | 17 ✓ |
| GGTTAA (-218, -213 bp) | rbcS-3A binding site consensus | 14 |
| | box II core rbcS-3A | 14 |
| TATTTTGCATAGATTACTCGTTTGACATTGTTGTTT (-737, -701 bp) | Similarity to | |
| TATAA (-28, -32 bp) | ORF1 <i>Drosophila</i> transposon | 18 |
| | putative TATA box | |

* complementary strand.






Fig. 2. Activity of the α -thionin promoter fused to the β -glucuronidase (GUS) gene in transgenic tobacco. (A) Representation of the gene fusion inserted in the Bin19 vector. (B) GUS activity in different parts of tobacco transformants carrying the α -TH-GUS construction (TH) and the CaMV-35S-GUS construction (35), and of non-transformed tobacco (N). Activity was determined fluorimetrically on a fresh weight basis [12] and protein was determined using the BCA kit (Pierce). Average yield of protein was 5.2 mg/g in seeds, 1.1 mg in leaves, 2.8 mg/g in stems, and 2.2 mg/g in roots. 1 unit = 1 pmol of 4-methylumbelliferone produced per min. (C) Temporal GUS expression in hand-dissected endosperms of α -TH-GUS (●), 35S-GUS (▲) and non-transformed (△) tobacco plants.

dosperm and reached a maximum at about 12 days after pollination, preceding by about four days the peak of the 35S control (Fig. 2C). This is in agreement with the expression of the α -thionin gene in barley [5], which also takes place in the first half of the development period.

Technical assistance from D. Lamoneda and J. García is acknowledged. Work was supported by a grant from the Fundación Ramón Areces.

References

- García-Olmedo, F., Rodríguez-Palenzuela, P., Hernández-Lucas, C., Ponz, F., Marañón, C., Carmona, M.J., López-Fando, J., Fernández, J.A. and Carbonero, P. (1989) *Plant Mol. Cell Biol.* 6, 31–60.
- García-Olmedo, F., Carmona, M.J., López-Fando, J.J., Fernández, J.A., Castagnaro, A., Molina, A., Hernández-Lucas, C. and Carbonero, P. (1991) in *Plant Gene Research Series: Genes Involved in Plant Defense* (Boller, T.H. and Meins, F., eds.), pp. 283–302, Springer Verlag, Berlin.
- Ponz, F., Paz-Ares, J., Hernández-Lucas, C., García-Olmedo, F. and Carbonero, P. (1986) *Eur. J. Biochem.* 156, 131–135.
- Hernández-Lucas, C., Royo, J., Paz-Ares, J., Ponz, F., García-Olmedo, F. and Carbonero, P. (1986) *FEBS Lett.* 200, 103–105.
- Rodríguez-Palenzuela, P., Pintor-Toro, J.A., Carbonero, P. and García-Olmedo, F. (1988) *Gene* 70, 271–281.
- Castagnaro, A., Marañón, C., Carbonero, P. and García-Olmedo, F. (1992) *J. Mol. Biol.* 224, 1003–1009.
- Bohlmann, H. and Apel, K. (1987) *Mol. Gen. Genet.* 207, 446–454.
- Bohlmann, H., Clausen, S., Behnke, S., Giese, H., Hiller, C., Reiman-Philipp, U., Schrader, G., Barkholt, V. and Apel, K. (1988) *EMBO J.* 7, 1559–1565.
- Gausling, K. (1987) *Planta* 171, 241–246.
- Fernández de Caley, R., Hernández-Lucas, C., Carbonero, P. and García-Olmedo, F. (1976) *Genetics* 83, 687–699.
- Bevan, M.W. (1984) *Nucleic Acid. Res.* 12, 8711–8721.
- Jefferson, R.A. (1987) *Plant Mol. Biol. Rep.* 5, 387–405.
- Castresana, C., García-Luque, I., Alonso, E., Malik, V.S. and Cashmore, A.R. (1988) *EMBO J.* 7, 1929–1936.
- Kuhlemeier, C., Fluhr, R., Green, P.J. and Chua, N.-H. (1987) *Genes Dev.* 1, 247–255.
- Lawton, M.A., Dean, S.M., Dron, M., Kooter, K.M., Kragh, K.M., Harrison, M.J., Yu, L., Tanguay, L., Dixon, R.A. and Lamb, C.J. (1991) *Plant Mol. Biol.* 16, 235–249.
- Bustos, M., Begum, D., Kalkan, F.A., Battraw, M.J. and Hall, T.C. (1991) *EMBO J.* 10, 1469–1479.
- Mark, S.T. and Richard, B.F. (1990) *Plant Cell* 2, 1171–1180.
- Templeton, N.S. and Potter, S.S. (1989) *EMBO J.* 8, 1887–1894.